An accuracy study for the DINAMAP SuperSTAT NIBP technology for adult/pediatric population

Andrea Stebor, RN, Ph.D., Rana F. Mikhail, Ph.D.

INTRODUCTION

The purpose of this clinical study was to determine the accuracy of the DINAMAP* SuperSTAT blood pressure technology in adult/ pediatric subjects according to the ANSI/AAMI SP-10:2002 Standard. In adults and children, SuperSTAT NIBP determinations acquired concurrently with the ECG parameter will report a determination in the presence of an irregular heart rhythm. Systolic, diastolic, mean arterial pressure (MAP), and pulse rate (PR) were compared to intraarterial blood pressure (IBP). The studies took place in cardiac catheterization labs. Institutional Review Board (IRB) approvals were granted and informed consent was obtained.

METHOD

Data were collected according to the ANSI/AAMI SP-10:2002 Standard specifications, utilizing static and dynamic calibration procedures to ensure the accuracy of the IBP. A computer program recorded the SuperSTAT NIBP simultaneously with the IBP. A sample of 28 adult/pediatric subjects (minimum age 4.02 years, maximum age 80.1 years) was in the data analysis. Four sizes of CRITIKON* DURACUF* blood pressure cuffs were used, child, small adult, adult, and large adult. Five subjects had atrial fibrillation. This was demonstrated by an irregular rhythm, inconsistent R-R interval, and absence of P waves on their ECG waveforms from the reference monitor in the cardiac catheterization unit. Every NIBP determination from these five subjects was reported in the presence of an irregular heart rhythm.

RESULTS

The differences between the SuperSTAT algorithm and the IBP reference were analyzed according to the ANSI/ AAMI SP-10:2002 Standard. *see* Table 1.

Parameter	Mean	Standard Deviation
Systolic difference (mmHg)	-0.451	4.448
Diastolic difference (mmHg)	-2.117	5.073
MAP difference (mmHg)	-1.773	3.833
Pulse rate difference (bpm)	-0.217	3.051

Table 1. Mean differences between SuperSTAT NIBP and the IBP reference:

CONCLUSION

The SuperSTAT NIBP algorithm met the accuracy specifications of \pm 5 mmHg for mean difference and \leq 8 mmHg standard deviation of the ANSI/AAMI SP-10:2002 Standard for the adult/pediatric population.

ADDITIONAL RESOURCES

For white papers, guides and other instructive materials about GE Healthcare's clinical measurements, technologies and applications, please visit http://clinicalview.gehealthcare.com/



Imagination at work

www.gehealthcare.com

©2014 General Electric Company – All rights reserved. GE and GE Monogram are trademarks of General Electric Company. GE Medical Systems Information Technology, a General Electric company, doing business as GE Healthcare. General Electric Company reserves the right to make changes in specifications and features shown herein, or discontinue the product described at any time without notice or obligation. Contact your GE Representative for the most current information. *Trademarks of General Electric Company.

JB25637XX 11/14